

Entire periodic functions with plane zeros

L.I.Ronkin

A.M.Russakovskii

Abstract

We give a complete description of divisors of entire periodic functions in $\mathbb{C}^n$ with plane zeros.

1 Introduction

The present note is close by the considered problems to the papers [Ro1, Ro2] and continues investigations of entire periodic functions of several complex variables and their divisors started in these papers.

In [Ro1] conditions were studied, under which an n - periodic positive divisor in $\mathbb{C}^n$ is a divisor of some entire n - periodic function. In [Ro2] a description was given, of those polynomials in n variables whose divisors after their n - periodic "reproduction" generate n - periodic divisors in $\mathbb{C}^n$. Here we give a complete description of divisors of entire periodic functions with plane zeros¹. Besides that we give a more thorough analysis of the conditions on divisors introduced in [Ro1].

We denote further by Z an n - periodic divisor with linearly independent periods $\omega_1 \in \mathbb{R}^n, \dots, \omega_n \in \mathbb{R}^n$ and by $f(z)$ an arbitrary entire function in $\mathbb{C}^n$ whose divisor Z_f coincides with Z . In this situation,

$$f(z + \omega_p) = e^{g_{\omega_p}(z)} f(z), \quad p = 1, \dots, n,$$

where the entire functions $g_p = g_{\omega_p}$ are defined by f up to constants of the form $2\pi im$, $m \in \mathbb{Z}$.

Denote by $\Delta_p = \Delta_{\omega_p}$ the difference operator

$$\Delta_p h = h(z + \omega_p) - h(z).$$

The integer quantities

$$N_{pq} = N(\omega_p, \omega_q : Z) \stackrel{\text{def}}{=} \frac{1}{2\pi i} (\Delta_p g_q - \Delta_q g_p)$$

¹The term "function with plane zeros" is utilized for functions whose zero set is a union of hyperplanes. Such functions have been object of study in a number of papers, particularly in [Gr, Se, Pa, PaRu]

do not depend on z nor on the choice of $f(z)$ (see next section). The skew-symmetric matrix

$$\widetilde{N}_Z = \{N_{pq}\}_{p,q=1}^n$$

will be called the *index* of the periodic² divisor Z .

Without loss of generality we assume henceforth that

$$\omega_1 = (1, 0, \dots, 0), \dots, \omega_n = (0, \dots, 0, 1).$$

Using the concept of the index of a periodic divisor, one can reformulate theorem 1 from [Ro1] in the following way.

Theorem A. *A periodic divisor $Z \subset \mathbb{C}^n$ is a divisor of some entire periodic function if and only if*

$$\widetilde{N}_Z = 0.$$

In [Ro1], a geometric sufficient condition for a divisor to be a divisor of a periodic entire function was also given:

Theorem B. *Let α_{pq} be the automorphism of $\mathbb{C}^n$ permuting the z_p and z_q coordinates.*

If a periodic divisor $Z \subset \mathbb{C}^n$ is symmetric, i.e. invariant with respect to α_{pq} , $\forall p, q$, then there exists an entire periodic function $F(z)$ with $Z_F = Z$.

In the next section, we make some elementary analysis of the properties of an index of a periodic divisor which allows to give a somewhat more general sufficient condition.

Theorem 1 *Let β_j be the automorphism of $\mathbb{C}^n$ taking z_j to $-z_j$ and not changing the rest of coordinates and let α_{pq} be the same as in theorem B.*

If for some splitting of the set $\{1, \dots, n\}$ into nonintersecting parts I and J a divisor $Z \subset \mathbb{C}^n$ is invariant with respect to β_k , $\forall k \in I$, and with respect to α_{pq} , $\forall p, q \in J$, then there exists an entire periodic function $F(z)$ with $Z_F = Z$.

The main subject of our study is periodic divisors with plane components, i.e. periodic divisors in $\mathbb{C}^n$ whose carrier is a union of hyperplanes. To formulate the main result of the paper, introduce additional denotations.

Let

$$l(z) = l_{a,c}(z) = \langle a, z \rangle + c, \quad a \in \mathbb{C}^n, \quad c \in \mathbb{C}$$

²For brevity, in what follows we will write simply "periodic divisor" instead of " n - periodic divisor with periods $\omega_1, \dots, \omega_n \in \mathbb{R}^n$ ".

and let

$$L = L_{a,c} = \{z \in \mathbb{C}^n : l(z) = 0\}$$

be the corresponding hyperplane. Denote by $\hat{S}_L$ the periodic "reproduction" of the hyperplane L , i.e.

$$\hat{S}_L = \bigcup_{k \in \mathbb{Z}^n} \{z \in \mathbb{C}^n : l(z - k) = 0\}.$$

Note that this operation does not necessarily produce an analytic set. If, however, it is analytic, we identify it with the divisor having $\hat{S}_L$ as its carrier with multiplicity 1 on it.

Consider the set

$$A_l = \{x \in \mathbb{R}^n : \langle x, a \rangle = 0\}.$$

This is a subspace in $\mathbb{R}^n$. Denote by m the dimension of its orthogonal complement in $\mathbb{R}^n$. Obviously, either $m = 1$ or $m = 2$. Denote the set of those linear functions $l(z)$ for which $m = 1$ by $\mathcal{L}_1$ and the set of those ones for which $m = 2$ by $\mathcal{L}_2$.

If $l_{a,c}$ is given, consider a "parallelogram"

$$P_{pq} = \{w \in \mathbb{C} : w = \alpha a_p + \beta a_q; 0 \leq \alpha < 1, 0 \leq \beta < 1\}.$$

If

$$a_q \neq 0 \text{ and } \operatorname{Im} \frac{a_p}{a_q} = 0,$$

then P_{pq} has no interior and becomes an interval.

Denote by ν_{pq} the number of points $w \in \mathbb{C}$ of the form $w = \langle k, a \rangle$, $k \in \mathbb{Z}^n$ belonging to P_{pq} . If a_p or a_q equals 0, we set $\nu_{pq} = 0$.

If $l \in \mathcal{L}_2$ and $\hat{S}_L$ is a divisor, then, as we will see in section 3,

$$\operatorname{Im} \frac{a_p}{a_q} \neq 0,$$

(and hence P_{pq} is nondegenerate) for at least one pair (p, q) .

Theorem 2 *A periodic divisor $Z \subset \mathbb{C}^n$ is a divisor of some entire periodic function with plane zeros if and only if it may be represented in the form*

$$Z = Z' + Z'',$$

where

$$Z' = \sum_{j=1}^{\mu_1} \hat{S}_{L_j}, \quad \mu_1 \leq \infty, \quad l_j = \langle a^{(j)}, z \rangle + c_j, \quad l_j \in \mathcal{L}_1$$

and

$$\lim_{j \rightarrow \infty} \frac{|\operatorname{Im} c_j|}{|a^{(j)}|} = \infty, \quad \text{if } \mu_1 = \infty,$$

and where

$$Z'' = \sum_{j=1}^{\mu_2} \hat{S}_{L_j}, \quad \mu_2 < \infty, \quad l_j = \langle a^{(j)}, z \rangle + c_j, \quad l_j \in \mathcal{L}_2$$

and

$$\sum_{j=1}^{\mu_2} \nu_{pq}^{(j)} \operatorname{sign} \operatorname{Im} \frac{a_q^{(j)}}{a_p^{(j)}}, \quad \forall p, q \text{ such that } a_p^{(j)} \neq 0.$$

Remark. In section 3 where this theorem is proved we give also the explicit form of the corresponding entire function.

2 Properties of the index and proof of theorem 1

In [Ro1] it was noted that the quantity N_{pq} does not depend on z and is an integer. It is easy to see also that this quantity does not depend on the choice of the function f defining the divisor Z . Indeed, if f and $\tilde{f}$ are entire functions with

$$Z_f = Z_{\tilde{f}} = Z,$$

then

$$f = \tilde{f} e^h$$

with h entire, and hence

$$g_p = \tilde{g}_p + \Delta_p h.$$

The function $\tilde{g}_p$ here is defined similar to g , i.e. by the equality

$$\tilde{f}(z + \omega_p) = \tilde{f}(z) e^{\tilde{g}_p(z)}.$$

We have then

$$\Delta_p g_q - \Delta_p \tilde{g}_q = \Delta_p \Delta_q h,$$

and due to commutativity of the operators Δ_p and Δ_q it follows that

$$\Delta_p g_q - \Delta_q g_p = \Delta_p \tilde{g}_q - \Delta_q \tilde{g}_p.$$

We outline below several elementary properties of the matrix $\tilde{N}_Z$.

1) if $k \in \mathbb{Z}$, $k > 0$, then

$$N(k\omega_p, \omega_q : Z) = kN(\omega_p, \omega_q : Z).$$

According to the definition, we have to determine functions $g_q(z) = g_{\omega_q}(z)$ and $g_p(z) = g_{k\omega_p}(z)$ first. If $f(z)$ is an entire function with $Z_f = Z$, then $g_p(z)$ is defined by the relation

$$f(z + k\omega_p) = f(z)e^{g_p(z)}.$$

Hence one easily sees that we may take

$$g_{k\omega_p}(z) = \sum_{j=0}^{k-1} g_{\omega_p}(z + j\omega_p).$$

Similarly, the operator $\Delta_p = \Delta_{k\omega_p}$ applied to a function h gives

$$[\Delta_{k\omega_p} h](z) = \sum_{j=0}^{k-1} [\Delta_{\omega_p} h](z + j\omega_p).$$

Therefore we have

$$\begin{aligned} 2\pi i N(k\omega_p, \omega_q : Z) &= \Delta_{k\omega_p} g_q - \Delta_{\omega_q} g_p \\ &= \sum_{j=0}^{k-1} \left[[\Delta_{\omega_p} g_q](z + j\omega_p) - [\Delta_{\omega_p} g_{\omega_p}](z + j\omega_p) \right] \\ &= k \cdot 2\pi i N(\omega_p, \omega_q : Z). \end{aligned}$$

2) If a divisor Z^* is obtained from the divisor Z with the help of the mapping

$$\beta_p : z \mapsto (z_1, \dots, z_{p-1}, -z_p, z_{p+1}, \dots, z_n),$$

then

$$N(\omega_p, \omega_q : Z^*) = -N(\omega_p, \omega_q : Z).$$

To show that this is true, note first that if $Z = Z_f$ for some entire function f then $Z^* = Z(f^*)$, where $f^*(z) = f(\beta_p z)$. Therefore the corresponding functions g_j^* for $j \neq p$ are related to g_j by the equalities $g_j^* = g_j(\beta_p z)$. For $j = p$ one has

$$e^{g_p^*(z)} = \frac{f^*(z + \omega_p)}{f^*(z)} = \frac{f(\beta_p z + \beta_p \omega_p)}{f(\beta_p z)} = \frac{f(\beta_p z - \omega_p)}{f(\beta_p z)} = e^{-g_p(\beta_p z - \omega_p)}$$

and hence one can take

$$g_p^*(z) = -g_p(\beta_p z - \omega_p).$$

From the above relations between g_j^* and g_j it follows that

$$\begin{aligned} 2\pi i N(\omega_p, \omega_q : Z^*) &= \Delta_p g_q^* - \Delta_q g_p^* \\ &= [g_q(\beta_p z + \beta_p \omega_p) - g_q(\beta_p z)] - [-g_p(\beta_p z - \omega_p + \omega_q) + g(\beta_p z - \omega_p)] \\ &= -[g_q(\beta_p z) - g_q(\beta_p z - \omega_p)] + [g_p(\beta_p z - \omega_p + \omega_q) - g(\beta_p z - \omega_p)] \\ &= (\Delta_q g_p - \Delta_p g_q)|_{\beta_p z - \omega_p} = -2\pi i N(\omega_p, \omega_q : Z). \end{aligned}$$

3) If a divisor Z^* is obtained from the divisor Z with the help of the mapping $\alpha_{pq} : \mathbb{C}^n \rightarrow \mathbb{C}^n$ permuting the z_p and z_q coordinates and leaving the other coordinates unchanged, then

$$N(\omega_p, \omega_q : Z^*) = -N(\omega_p, \omega_q : Z).$$

To prove this, consider as before functions f, g_p and f^*, g_p^* corresponding to the divisors Z and Z^* . Evidently, one can take $f^*(z) = f(\alpha_{pq}(z))$. Then

$$\frac{f^*(z + \omega_p)}{f^*(z)} = \frac{f(\alpha_{pq}z + \alpha_{pq}\omega_p)}{f(\alpha_{pq}z)} = \frac{f(\alpha_{pq}z + \omega_q)}{f(\alpha_{pq}z)},$$

whence one can take

$$g_p^* = g_q(\alpha_{pq}z).$$

This implies

$$\begin{aligned} 2\pi i N(\omega_p, \omega_q : Z^*) &= [g_q^*(z + \omega_p) - g_q^*(z)] - [g_p^*(z + \omega_q) - g_p^*(z)] \\ &= [g_p(\alpha_{pq}z + \omega_q) - g_p(\alpha_{pq}z)] - [g_q(\alpha_{pq}z + \omega_p) - g_q(\alpha_{pq}z)] \\ &= (\Delta_q g_p - \Delta_p g_q)|_{\alpha_{pq}z} = 2\pi i N(\omega_q, \omega_p : Z) = -2\pi i N(\omega_p, \omega_q : Z) \end{aligned}$$

4) If a divisor Z is a sum of periodic divisors Z_1 and Z_2 then

$$\widetilde{N}_Z = \widetilde{N}_{Z_1} + \widetilde{N}_{Z_2}.$$

This property is obvious.

We would like to note that the theorem B mentioned in the introduction follows immediately from theorem A and the property 3). In [Ro1] it was proved (also deduced from theorem A) in a less elementary way.

If one uses the property 2) together with the property 3), then theorem A yields immediately our theorem 1.

3 Linear periodic divisors. Proof of theorem 2

Next we consider periodic divisors with plane components, i.e. divisors whose carrier is the union of hyperplanes of the form $\{z : \langle a, z \rangle + c = 0\}$.

Let, as before,

$$l(z) = l_{a,c}(z) = \langle a, z \rangle + c, \quad a \in \mathbb{C}^n, \quad c \in \mathbb{C}$$

and let

$$A = A_l = \{x \in \mathbb{R}^n : l(z - x) \equiv l(z)\} = \{x \in \mathbb{R}^n : \langle a, x \rangle = 0\},$$

$$A^\perp = A_l^\perp = \{\xi \in \mathbb{R}^n : \langle \xi, x \rangle = 0, \forall x \in A\}.$$

Denote the dimension of $A^\perp$ by m . Let $\Lambda_1, \dots, \Lambda_m$ be a basis in $A^\perp$ and $\Lambda_{m+1}, \dots, \Lambda_n$ be a basis in A . Then the vectors $\Lambda_1, \dots, \Lambda_n$ form a basis in both $\mathbb{R}^n$ and $\mathbb{C}^n$ (over $\mathbb{R}$ and $\mathbb{C}$ respectively). Hence one can represent $z \in \mathbb{C}^n$ as follows:

$$z = \sum_{j=1}^n \zeta_j(z) \Lambda_j,$$

where $\zeta_j(z) = \langle \Omega_j, z \rangle$ and Ω_j is the j -th row of the matrix Ω reciprocal to the matrix Λ with columns $\Lambda_1, \dots, \Lambda_n$.

From the definition of the spaces $A^\perp$ and A it follows that

$$l(z) = c + \langle a, \zeta_1(z) \Lambda_1 + \dots + \zeta_m(z) \Lambda_m \rangle = l^*(\zeta')|_{\zeta'=\zeta'(z)},$$

where

$$\zeta' = (\zeta_1, \dots, \zeta_m), \quad l^*(\zeta') = \sum_{j=1}^m b_j \zeta_j, \quad b_j = \langle a, \Lambda_j \rangle.$$

Let L be the hyperplane $\{z : l(z) = 0\}$. Consider the periodic "reproduction" of the hyperplane L , i.e.

$$\hat{S}_L = \bigcup_{k \in \mathbb{Z}^n} \{z \in \mathbb{C}^n : l(z - k) = 0\}.$$

This set is obviously periodic and if it is analytic, it can be considered as a periodic divisor, i.e. the carrier of the divisor is $\hat{S}_L$ and the multiplicity on it equals 1.

From the results in [Ro2] it follows in particular that $\hat{S}_L$ will be a divisor if and only if the space $A^\perp$ possesses a basis consisting of vectors with integer coordinates and the function $l^*(\zeta')$ is hypoelliptic, that is, satisfies the condition: *the distance from a point $\xi' \in \mathbb{R}^m$ to the set $\{\zeta' \in \mathbb{C}^m : l^*(\zeta') = 0\}$ tends to ∞ when ξ' tends to ∞ .*

Let $\widehat{S}_L$ be a divisor. We are interested in the question when this divisor is a divisor of some entire periodic function.

We have already split all linear functions $l(z)$ into two sets $\mathcal{L}_1$ and $\mathcal{L}_2$ with regard to the dimension m of the space $A^\perp$.

Consider the case $l \in \mathcal{L}_1$ first. In this case

$$A^\perp = \{x \in \mathbb{R}^n : x = ta^0, t \in \mathbb{R}\}, \quad a^0 \in \mathbb{R}^n, \quad a = w_0 a^0,$$

where $w_0 \in \mathbb{C}$, and if $\widehat{S}_L$ is a divisor, without loss of generality we may assume that $a = a^0 = k^0 \in \mathbb{Z}^n$, and that the coordinates of k^0 have no common divisor.

Put

$$f_l(z) = \sin[\pi(< k^0, z > + c)]e^{i\pi(< k^0, z > + c)}.$$

Evidently, f_l is an entire periodic function with divisor

$$Z_f = \bigcup_{j=-\infty}^{\infty} \{z : < k^0, z > + c = j\} = \bigcup_{k \in \mathbb{Z}^n} \{z : < k^0, z > + c = < k^0, k >\} = \widehat{S}_L.$$

Hence we get an affirmative answer to our question when $l \in \mathcal{L}_1$.

Let now $l \in \mathcal{L}_2$. Then

$$l^*(\zeta') = b_1 \zeta_1 + b_2 \zeta_2 + c,$$

and

$$l(z) = b_1 \zeta_1(z) + b_2 \zeta_2(z) + c,$$

where

$$b_1 = < a, \Lambda_1 >, \quad b_2 = < a, \Lambda_2 >, \quad \zeta_1(z) = < \Omega_1, z >, \quad \zeta_2(z) = < \Omega_2, z >.$$

If $\widehat{S}_L$ is a divisor, then the function $l^*(\zeta')$ is hypoelliptic, and we may assume that the elements of the matrix Λ are integers. It is immediate to check that the function $l^*(\zeta_1, \zeta_2)$ is hypoelliptic if and only if

$$b_1 \neq 0, \quad b_2 \neq 0, \quad \text{and} \quad \text{Im} \frac{b_2}{b_1} \neq 0,$$

or equivalently, when $\text{Im} \frac{a_p}{a_q} \neq 0$ for at least one pair of indices p and q .

First we consider the simplest case with regard to the number of variables, n . Note that if $n = 2$, the original function $l(z_1, z_2)$ is also hypoelliptic. To compute the index of

the divisor $Z = \widehat{S}_L$ in this case, we construct an entire function of one complex variable w which has simple zeros precisely at points $w = p + qT$, $(p, q) \in \mathbb{Z}^2$ where $T = \frac{a_2}{a_1}$.

Suppose that $\text{Im } T > 0$ and consider the product

$$\Phi_T(w) = \prod_{q=0}^{\infty} \{2ie^{-i\pi(w-qT)} \sin \pi(w-qT)\} \prod_{q=-\infty}^{-1} \{2ie^{i\pi(w-qT)} \sin \pi(w-qT)\} \quad (1).$$

Since

$$2ie^{-iw} \sin w = 1 + O(e^{-2|\text{Im } w|}) \quad \text{when } \text{Im } w \rightarrow -\infty,$$

and

$$2ie^{iw} \sin w = 1 + O(e^{-2|\text{Im } w|}) \quad \text{when } \text{Im } w \rightarrow +\infty,$$

the above product converges and hence is an entire function.

Similarly, if $\text{Im } T < 0$ we set

$$\Phi_T(w) = \prod_{q=0}^{\infty} \{2ie^{i\pi(w-qT)} \sin \pi(w-qT)\} \prod_{q=-\infty}^{-1} \{2ie^{-i\pi(w-qT)} \sin \pi(w-qT)\} \quad (2).$$

Next we denote

$$u_q^{\pm}(w) = 2ie^{\pm i\pi(w-qT)} \sin \pi(w-qT).$$

It is obvious that

$$u_q^{\pm}(w+1) = u_q^{\pm}(w) \quad \text{and} \quad u_q^{\pm}(w+T) = u_{q-1}^{\pm}(w).$$

Hence

$$\Phi_T(w+1) = \Phi_T(w) \quad (3)$$

and for $\text{Im } T > 0$

$$\Phi_T(w+T) = \prod_{q=0}^{\infty} u_q^-(w+T) \prod_{q=-\infty}^{-1} u_q^+(w+T) = \frac{u_{-1}^-(w)}{u_{-1}^+(w)} \Phi_T(w) = e^{-2\pi i(w+T)} \Phi_T(w). \quad (4)$$

Similarly, when $\text{Im } T < 0$,

$$\Phi_T(w+T) = e^{2\pi i(w+T)} \Phi_T(w). \quad (5)$$

Consider the function

$$f(z_1, z_2) = f_{a,c}(z_1, z_2) = \Phi_T(z_1 + Tz_2 + \frac{c}{a_1}). \quad (6)$$

This entire function vanishes if and only if

$$z_1 + Tz_2 + \frac{c}{a_1} = p + qT, \quad (p, q) \in \mathbb{Z}^2,$$

that is, when

$$l(z_1 - p, z_2 - q) = 0, \quad (p, q) \in \mathbb{Z}^2.$$

Thus $Z_f = \widehat{S}_L$.

From (3)-(6) it follows that the corresponding functions g_1 and g_2 may be taken in the following form:

$$g_1 \equiv 0$$

and

$$g_2(z) = (-\text{sign Im } T) \cdot 2\pi i(z_1 + Tz_2 + \frac{c}{a_1} + T).$$

Therefore

$$N(\omega_1, \omega_2 : \widehat{S}_L) = -\text{sign Im } T,$$

and hence the divisor $\widehat{S}_L$ for $l \in \mathcal{L}_2$ cannot be a divisor of any entire periodic function.

Assume now that the dimension $n > 2$, and let $l \in \mathcal{L}_2$. Suppose $\text{Im } \frac{a_2}{a_q} \neq 0$ for some p and q .

Fix these p and q . To compute N_{pq} , we will construct a function $f(z)$ having divisor $Z_f = \widehat{S}_L$ in the form

$$f(z) = \Psi\left(\frac{l(z)}{a_p}\right),$$

where the entire function $\Psi(w)$, $w \in \mathbb{C}$, periodic with period 1, is to be defined below.

Let $T_j = \frac{a_j}{a_p}$, $j = 1, \dots, n$. In order that $Z_f = \widehat{S}_L$, zeros of the function $\Psi(w)$ must be simple and have the form

$$\{w \in \mathbb{C} : w = k_p + \sum_{j=1}^n k_j T_j\}_{k=(k_1, \dots, k_n) \in \mathbb{Z}^n},$$

i.e. they must make up a periodic set Z_Ψ with n periods $T_1, \dots, T_n$.

Denote by $\mathcal{A}_l$ the set

$$\mathcal{A}_l = \{w \in \mathbb{C}^n : w = \langle a, k \rangle, \quad k \in \mathbb{Z}^n\}$$

and let $x_1 = 0, \dots, x_{\nu_{pq}}$ be all the points from $\mathcal{A}_l$ belonging to the parallelogram

$$P_{pq} = \{w : w = \alpha a_p + \beta a_q, \quad 0 \leq \alpha < 1, 0 \leq \beta < 1\}.$$

Then

$$\mathcal{A}_l = \bigcup_{j=1}^{\nu_{pq}} A_j,$$

where

$$A_j = \{w \in \mathbb{C} : w = k_p a_p + k_q a_q + x_j, (k_p, k_q) \in \mathbb{Z}^2\}.$$

From all above it follows that Z_Ψ may be represented in the form of a union of ν_{pq} identical nonintersecting sets

$$Z_j = \{w \in \mathbb{C} : a_p w \in A_j\}$$

each having periods 1 and T_q , and being shifted relative to each other:

$$Z_\Psi = \bigcup_{j=1}^{\nu_{pq}} (Z_1 + \frac{x_j}{a_p}).$$

Using the functions $\Phi_T(w)$ constructed above, we set

$$\Psi(w) = \bigcup_{j=1}^{\nu_{pq}} \Phi_{T_q}(w - \frac{x_j}{a_p}).$$

By virtue of our construction the function

$$f(z) = \Psi(\frac{l(z)}{a_p})$$

has divisor $\widehat{S}_L$.

Since the function Ψ is periodic with period 1, one can take

$$g_p(z) = \log \frac{f(z + \omega_p)}{f(z)} \equiv 0,$$

and

$$g_q(z) = -\text{sign Im } T_q \cdot 2\pi i \sum_{j=1}^{\nu_{pq}} (\frac{l(z)}{a_p} - \frac{x_j}{a_p} + T_q), \quad (7)$$

which implies that

$$N(\omega_p, \omega_q : \widehat{S}_L) = -\nu_{pq} \cdot \text{sign Im } T_q \neq 0.$$

Therefore the divisor $\widehat{S}_L$ cannot be a divisor of any entire periodic function.

Let, as above, $l \in \mathcal{L}_2$ and assume that $\widehat{S}_L$ is a divisor. In what follows we will need to compute N_{pq} also for such p and q that $\text{Im } \frac{a_p}{a_q} = 0$. Note that in this case $\frac{a_q}{a_p}$ is a rational number since in the two-dimensional (over $\mathbb{Z}$) set $\{\frac{a_j}{a_p}, j = 1, \dots, n\}$ there are already two $\mathbb{Z}$ -independent periods (1 and $\frac{a_i}{a_p} \notin \mathbb{R}$ for some i). So, there are two integers, k and j such that

$$ja_p = ka_q.$$

To compute N_{pq} we use properties 1) and 2) of the index. In view of these properties

$$N(\omega_p, \omega_q : Z) = \frac{1}{k} N(\omega_p, k\omega_q : Z),$$

so it is enough to compute $N(\omega_p, k\omega_q : Z)$. By the above definition of $f(z)$,

$$e^{g_{k\omega_q}(z)} = \frac{f(z + k\omega_q)}{f(z)} = \frac{\Psi(\frac{l(z+k\omega_q)}{a_p})}{\Psi(\frac{l(z)}{a_p})} = \frac{\Psi(\frac{l(z)}{a_p} + \frac{ka_q}{a_p})}{\Psi(\frac{l(z)}{a_p})} = \frac{\Psi(\frac{l(z)}{a_p} + j)}{\Psi(\frac{l(z)}{a_p})} = 1,$$

since the function $\Psi(z)$ has period 1 and $j \in \mathbb{Z}$. Therefore we can take $g_{k\omega_q} \equiv 0$ and since $g_p = g_{\omega_p} \equiv 0$, it follows that $N(\omega_p, k\omega_q : Z) = 0$, and consequently $N_{pq} = 0$.

Now we have all the necessary information to describe the structure of divisors Z of entire periodic functions with plane zeros. Such a divisor is obviously periodic and has only hyperplane irreducible components. From the above considerations it follows that a periodic divisor with plane components may be represented in the form

$$Z = Z' + Z'',$$

where

$$Z' = \sum_{j=1}^{\mu_1} \widehat{S}_{L_j}, \quad l_j \in \mathcal{L}_1, \quad \mu_1 \leq \infty,$$

and

$$Z'' = \sum_{j=1}^{\mu_2} \widehat{S}_{L_j}, \quad l_j \in \mathcal{L}_2, \quad \mu_2 \leq \infty,$$

so that each $\widehat{S}_{L_j}$ is a divisor itself.

It is immediate to check that each function $l \in \mathcal{L}_2$ vanishes at some point of $\mathbb{R}^n = \mathbb{R}^n + i0$. Hence the corresponding set $\widehat{S}_L$ has nonempty intersection with the cube

$$\{x \in \mathbb{R}^n : |x_j| \leq 1, j = 1, \dots, n\}.$$

Since a divisor can have only a finite number of irreducible components intersecting a fixed compact, it follows that $\mu_2 < \infty$ and hence

$$\widetilde{N}_{Z''} = \sum_{j=1}^{\mu_2} \widetilde{N}_{Z_j},$$

where $Z_j = \widehat{S}_{L_j}$.

From theorem A and property 4) of indices it follows now that the divisor Z'' will be a divisor of an entire periodic function if and only if

$$\sum_{j=1}^{\mu_2} N(\omega_p, \omega_q : \widehat{S}_{L_j}) = 0, \quad \forall p, q.$$

In terms of coefficients ($l_j = \langle a^{(j)}, z \rangle + c_j$) the last condition may be rewritten as

$$\sum_{j=1}^{\mu_2} \nu_{pq}^{(j)} \operatorname{sign} \left(\operatorname{Im} \frac{a_q^{(j)}}{a_p^{(j)}} \right) = 0, \quad \forall p, q = 1, \dots, n \text{ with } a_p^{(j)} \neq 0. \quad (8)$$

The corresponding entire periodic function, i.e. entire periodic function F_2 with $Z_{F_2} = Z''$ is constructed as follows.

First, we make additional investigation of the case

$$l \in \mathcal{L}_2, \widehat{S}_L \text{ is a divisor.}$$

As it has been noted before, in this case

$$l(z) = b_1 \zeta_1(z) + b_2 \zeta_2(z),$$

where

$$b_1 \neq 0, b_2 \neq 0, \operatorname{Im} \frac{b_2}{b_1} \neq 0, \zeta_1 = \langle k, \Omega_1 \rangle, \zeta_2 = \langle k, \Omega_2 \rangle,$$

and Ω_1, Ω_2 are vectors with rational coordinates. The set

$$\mathcal{A}_l \stackrel{\text{def}}{=} \{w = \langle a, k \rangle, k \in \mathbb{Z}^n\}$$

has the form

$$\mathcal{A}_l = \{w : w = \langle k, b_1 \Omega_1 + b_2 \Omega_2 \rangle\}.$$

Since Ω_1 and Ω_2 have rational coordinates, the set $\mathcal{A}_l$, which is obviously an additive group, has no density points in $\mathbb{C}$ and is therefore a free group with 2 generators. Hence there exist such

$$w_1 = w_1(l) \in \mathbb{C} \setminus 0, w_2 = w_2(l) \in \mathbb{C} \setminus 0, \frac{w_2}{w_1} = T(l) \in \mathbb{C} \setminus \mathbb{R},$$

that

$$\mathcal{A}_l = \{w : w = j_1 w_1 + j_2 w_2, (j_1, j_2) \in \mathbb{Z}^2\}.$$

Particularly,

$$a_p = m_{1p} w_1 + m_{2p} w_2, \quad m_{ip} = m_{ip}(l) \in \mathbb{Z}, \quad i = 1, 2.$$

Set

$$F_{(l)}(z) = \Phi_T\left(\frac{l(z)}{w_1}\right),$$

where the function Φ is defined by (1) or (2) depending on the sign of $\text{Im } T(=T(l))$.

Then

$$Z_{F_{(l)}} = \bigcup_{(j_1, j_2) \in \mathbb{Z}^2} \{z : l(z) = j_1 w_1 + j_2 w_2\} = \bigcup_{k \in \mathbb{Z}^n} \{z : l(z) = \langle a, k \rangle\} = \widehat{S}_L,$$

and by (3) and (4)

$$F_{(l)}(z + \omega_p) = e^{\widetilde{g}_{p,l}(z)} F_{(l)}(z),$$

with

$$\widetilde{g}_{p,l}(z) = -2\pi i \text{sign } T(l) \left[\frac{m_{1p} l(z)}{w_1} + \frac{m_{1p}(m_{1p} + 1)}{2} T(l) \right].$$

Getting back to our divisor $Z'' = \bigcup_{j=1}^{\mu_2} \widehat{S}_{L_j}$, set

$$\widetilde{F} = \prod_{j=1}^{\mu_2} F_{(l_j)}(z). \tag{9}$$

From all above it follows that

$$Z_{\widetilde{F}} = Z''$$

and that

$$\widetilde{F}(z + \omega_p) = e^{G_p(z)} \widetilde{F}(z)$$

with

$$G_p(z) = 2\pi i \sum_{j=1}^{\mu_2} \left(\text{sign } \text{Im } T(l_j) \cdot \left[\frac{m_{1p}(l_j) l_j(z)}{w_1^{(j)}} + \frac{m_{1p}(l_j)(m_{1p}(l_j) + 1)}{2} T(l_j) \right] \right).$$

Therefore the functions $G_p(z)$ are linear in all variables, i.e. have the form

$$\sum_{j=1}^n \sigma_j^p z_j + \tau_p.$$

Since $N(\omega_p, \omega_q : Z'') = 0$ and since $\Delta_q G_p = \sigma_q^p$, it follows that $\sigma_q^p = \sigma_p^q$, $\forall p, q$.

Put

$$H(z) = \frac{1}{2} \sum_{j=1}^n \sigma_j^j z_j^2 + \sum_{j=1, j \neq p}^n \sigma_j^p z_j z_p + \sum_{j=1}^n (\tau_j - \frac{1}{2} \sigma_j^j) z_j. \quad (10)$$

Obviously,

$$\Delta_p H = G_p.$$

Therefore the function

$$F_2(z) = \tilde{F}(z) e^{-H(z)} \quad (11)$$

will be the required periodic function with divisor Z'' .

It remains to consider the sum Z' . This sum will be a divisor if and only if each compact in $\mathbb{C}^n$ is intersected only by a finite number of divisors $\hat{S}_{L_j}$. Since $l_j \in \mathcal{L}_1$, we may assume that each function l_j has the form

$$l_j = \langle k^{(j)}, z \rangle + c_j, \quad k^{(j)} \in \mathbb{Z}^n, \quad c_j \in \mathbb{C}.$$

Hence the projection of the divisor $\hat{S}_{L_j}$ onto $\mathbb{R}^n$ has nonempty intersection with each cube

$$\{x \in \mathbb{R}^n : t_j \leq x_j \leq t_j + 1, \quad j = 1, \dots, n\},$$

and the projection of $\hat{S}_{L_j}$ onto $i\mathbb{R}^n$ is the plane

$$\{iy : y \in \mathbb{R}^n, \quad \langle k^{(j)}, y \rangle + \text{Im } c_j = 0\}.$$

Therefore the sum $\sum_{j=1}^{\mu_1} \hat{S}_{L_j}$ is a divisor if and only if either $\mu_1 < \infty$ or $\mu_1 = \infty$ and

$$\lim_{j \rightarrow \infty} \frac{|\text{Im } c_j|}{|k^{(j)}|} = \infty.$$

It was shown above that each divisor $\hat{S}_{L_j}$ is a divisor of some entire periodic function. So if $\mu_1 < \infty$, the divisor Z' will also be a divisor of a periodic function equal to the product of the corresponding functions for $\hat{S}_{L_j}$. We will show that a similar fact holds also in the case $\mu_1 = \infty$.

In order to prove this set

$$\gamma_j = \frac{\text{Im } c_j}{|k^{(j)}|}$$

and form the product

$$F_1(z) = \prod_{j=1}^{\infty} \phi_j(z). \quad (12)$$

with

$$\phi_j = 2i \sin \pi(< k^{(j)}, z > + c_j) \cdot e^{\text{sign } \gamma_j i \pi (< k^{(j)}, z > + c_j)} \quad (13).$$

Taking into account the above mentioned asymptotics of $2ie^w \sin w$ we obtain that for $|\text{Im } z| < R$,

$$F_1(z) = \prod_{j=1}^{\infty} (1 + O(e^{-|k^{(j)}|(|\gamma_j| - R)})). \quad (14)$$

By our assumption, $|\gamma_j| \rightarrow \infty$. Hence (14) implies that the product F_1 converges. Summarizing all above we come to the following conclusion:

Theorem 2. *A periodic divisor $Z \subset \mathbb{C}^n$ is a divisor of some entire periodic function $F(z)$ with plane zeros if and only if it may be represented in the form*

$$Z = Z' + Z'',$$

where

$$Z' = \sum_{j=1}^{\mu_1} \hat{S}_{L_j}, \quad \mu_1 \leq \infty, \quad l_j = < a^{(j)}, z > + c_j, \quad l_j \in \mathcal{L}_1$$

and

$$\lim_{j \rightarrow \infty} \frac{|\text{Im } c_j|}{|a^{(j)}|} = \infty, \quad \text{if } \mu_1 = \infty,$$

and where

$$Z'' = \sum_{j=1}^{\mu_2} \hat{S}_{L_j}, \quad \mu_2 < \infty, \quad l_j = < a^{(j)}, z > + c_j, \quad l_j \in \mathcal{L}_2$$

and

$$\sum_{j=1}^{\mu_2} \nu_{pq}^{(j)} \text{sign } \text{Im } \frac{a_q^{(j)}}{a_p^{(j)}}, \quad \forall p, q \text{ such that } a_p^{(j)} \neq 0.$$

Under these conditions the function $F(z)$ is representable in the form

$$F(z) = e^{\Psi(z)} F_1(z) F_2(z),$$

where $\Psi(z)$ is an arbitrary entire periodic function, and the functions $F_1(z)$ and $F_2(z)$ are constructed with regard to divisors Z' and Z'' by the above mentioned method, i.e. by virtue of (9) - (14).

References

- [Gr] L.Gruman. The regularity of growth of entire functions whose zeros are hyperplanes. Ark. Mat. 10, 1972, p. 23-31.
- [Pa] D. E. Papush. Entire functions of several variables with a regular set of 'plane' zeros. Sib. Math. J., 32, 1991, p. 120-130.
- [PaRu] D.E.Papush, A.M.Russakovskii. Interpolation on plane sets in $\mathbb{C}^2$. Ann. Fac. Sci. Toulouse, 1, n 3, 1992, p. 337 - 362.
- [Ro1] L.I.Ronkin. Entire periodic functions of several variables, periodic divisors. To appear in Math. Physics, Math. Anal. and Geom.
- [Ro2] L.I.Ronkin. Periodic algebraic divisors in $\mathbb{C}^n$. To appear in Math. Physics, Math. Anal. and Geom.
- [Se] A.B.Sekerin. On construction of entire functions with given growth. Sib. Math. J., 27, n 3, 1986, p. 179 - 192.

Theory of Functions Department
Mathematical Division
Institute for Low Temperature Physics
47 Lenin Avenue
310164 Kharkov
Ukraine